

HYGIENE SWAB

CODE	DESCRIPTION
MW570125	Listeria Isolation Transwab® in peel pouch (125 pieces)
MW57025	Listeria Isolation Transwab® in peel pouch (25 pieces)

Intended Use

Listeria Isolation Transwab is designed to be used alongside traditional selective methods to improve the quality system and minimise the risk of Listeria contamination. This simple to use diagnostic test can be applied anywhere in the environment where the presence of *Listeria spp.* would be critical. *Listeria spp.* and specifically *Listeria monocytogenes* are rapidly becoming the most important pathogen in the Food Industry; regulatory bodies from around the world are insisting that all food products are Listeria free.

Listeria Isolation Transwab works on an enhanced esculin media formulation. The hydrolysis of esculin gives a distinctive black/brown precipitate. Inhibitors are present in the media which will inhibit the growth of non *Listeria spp.* Only *Listeria spp.* will grow within the media to produce a black/brown precipitate due to esculin hydrolysis (table 3)⁵.

Instructions for Use

1. Peel back wrapper to expose both caps.
2. Remove swab and sample test site by rubbing swab over as large an area as practicable.
3. Remove cap of culture tube with thumb and forefinger and discard.
4. Insert swab in culture tube and push down fully to immerse the swab completely.
5. Fill in time, date and site details.
6. Incubate at 37°C for up to 48 hours.
7. Read results. A positive result is indicated by a colour change from Light Brown Agar to Black/Dark Brown commencing around the bud, any colour change is significant. Negative result shows no colour change.
8. Record results and dispose of tube.

Precautions

If any black media tubes are found as soon as the box is opened DO NOT USE.

Storage

Store the devices at 5-25°C
Do not freeze.

Shelf Life

12 months from the date of manufacture. Expiration date is printed on the tube label, peel pouch and box label.

Precautions when swabbing

1. When sampling an area always use the same swabbing technique.
2. Always try to sample the same surface area (100cm² or 200cm²), and be consistent in your sampling.
3. The moisture level on the surface should not vary at different sample times. It may be preferable with dry surfaces to wet the swab in a suitable wetting agent such as Ringers, Tween Lecithin Thiosulphate Ringers or Tween Peptone water.



Battery of Organisms Tested against Listeria Isolation Transwab®

<u>Organism</u>	<u>C.F.U. in Sample</u>	<u>Colour Change</u>
<i>Listeria monocytogenes</i> NCTC 11994	7	Black
<i>Listeria monocytogenes</i> NCTC 5214	19	Black
<i>Listeria monocytogenes</i> NCTC 7973	15	Black
<i>Listeria ivanovii</i>	10	Black
<i>Listeria innocua</i> NCTC 11288	9	Black
<i>Listeria seeligeri</i>	1 x 10 ⁴	Black
<i>Listeria welshimeri</i> NCTC 11857	1 x 10 ³	Black
<i>Listeria murrayi</i>	1 x 10 ⁵	Black
<i>Listeria grayi</i>	1 x 10 ⁵	Black
<i>Staphylococcus aureus</i> NCTC 6571	4 x 10 ⁸	No Change



<i>Strep faecalis</i> NCTC 775	1.5 x 10 ⁵	No Change
<i>Aeromonas hydrophilla</i> NCTC 1767	1.5 x 10 ⁷	No Change
<i>Bacillus subtilis</i> NCTC 10400	4 x 10 ⁷	No Change
<i>Bacillus pumilis</i> NCTC 10327	1 x 10 ⁷	No change
<i>Bacilus cereus</i> NCTC 10320	2 x 10 ⁶	No change
<i>Escherichia coli</i> NCTC 9001	1 x 10 ⁷	No change
<i>Escherichia coli</i> NCTC 10418	6 x 10 ⁷	No change
<i>Klebsiella pneumoniae</i>	3 x 10 ⁷	No change
<i>Klebsiella aerogenes</i> NCTC 7418	1 x 10 ⁸	No change
<i>Klebsiella aerogenes</i> NCTC 11228	3 x 10 ⁷	No change
<i>Proteus vulgaris</i> NCTC 1683	1 x 10 ⁸	No change
<i>Proteus mirabilis</i> NCTC 841	5 x 10 ⁸	No change
<i>Citrobacter freundii</i>	1.2 x 10 ⁸	No change
<i>Citrobacter diversus</i>	3 x 10 ⁶	No change
<i>Enterobacter agglomerans</i>	1.1 x 10 ⁸	No Change
<i>Morganella morgani</i>	1.4 x 10 ⁸	No Change
<i>Serratia liquefaciens</i>	1 x 10 ⁸	No Change
<i>Serratia marcescens</i>	1 x 10 ⁷	No Change
<i>Yersinia enterolytica</i>	1 x 10 ⁷	No Change
<i>Enterobacter cloacae</i>	3 x 10 ⁷	No Change
<i>Salmonella typhimurium</i> NCTC 74	3 x 10 ⁸	No Change
<i>Shigella flexneri</i>	8 x 10 ⁶	No Change
<i>Psuedomonas aeruginosa</i>	9 x 10 ⁷	No Change
<i>Pseudomonas putida</i> NCTC 10936	1.5 x 10 ⁷	No Change
<i>Pseudomonas fluorescens</i> NCTC 10038	1 x 10 ⁷	No Change
<i>Pseudomonas maltophilia</i> NCTC 102157	6 x 10 ⁷	No Change
<i>Pseudomonas vesicularis</i> NCTC 10900	4 x 10 ⁶	No Change
<i>Pseudomonas cepacia</i> NCTC 10743	3.5 x 10 ⁷	No Change
<i>Pseudomonas putrefaciens</i> NCTC 10735	3 x 10 ⁶	No Change
<i>Candida albicans</i>	2 x 10 ⁶	No Change

Example of area to be swabbed



STERILE R



10cm

